



كلية الطب
جامعة النهرين
تأسست عام 1987

AL-Nahrain University

College of Medicine

Department of Surgery

**Functional results after partial arthroplasty in
fracture neck femur in elderly patients**

A PROGECT

SUBMITTED TO THE AL-NAHRAIN COLLEGE OF
MEDICINE/DEPARTMENT OF SURGERY/ORTHOPAEDIC
SURGERY IN PARTIAL FULLFILMENT OF THE
REQUIRMENT OF THE BACHELOR DEGREE IN GENERAL
MEDICINE AND SURGERY

BY

Medical student

Ban Dawood Saleem

Supervisor

Assistant Professor Dr.Ahmed Sabeeh

M.B.Ch.B, F.I.B.M.S (Orthopedic Surgeon)

2019 A.D.

1440 AH

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

((يرفع الله الذين آمنوا منكم والذين أوتوا العلم درجات
والله بما تعلمون خبير))

صدق الله العظيم

((المجادلة 11))

Dedication

I dedicate this work to my beloved family, and to whom who support me.

Acknowledgement

First of all, most of all thanks go to the grace of God, for giving me the power and the intention to accomplish this work.

I would like to express my deepest appreciation and utmost gratitude to my supervisor, Dr.Ahmed Sabeeh for his encouragement, guidance, effort, interest and patience.

My special gratitude is to whom who support me to finish my study

SUMMARY

Background: fracture of neck femur is common in clinical practice. When elderly people are taken into consideration and the poor outcome in the form of morbidity and mortality in association with these fractures is also frequent.

Aim of this study: To evaluate the functional results after partial arthroplasty in compares to others modalities of management of fracture neck femur in elderly patients.

Literatures' review:

Femoral neck fractures by themselves are challenging to orthopedic surgeon and become more challenging because of the high rate of medical comorbidities in such elderly patients. It is well known that surgical intervention is the best approach in such fractures; however, the most fruitful surgical option is still controversial. The patient with non-displaced femoral neck fracture especially elderly one with chronic medical conditions can be treated with nonsurgical method. Others patients can be treated by surgical fixation with complications such as nonunion and osteonecrosis that could occur.

Displaced femoral neck fracture is at significant risk of non- union and osteonecrosis. Treatment option include closed reduction or ORIF(open reduction-internal fixation) with different construct, hemiarthroplasty(unipolar and bipolar)and total hip arthroplasty THA .Internal fixation of femoral neck fracture is associated with a greater number of significant problems(e.g. osteonecrosis,nonunion,revision)over hemiarthroplasty. Cemented prostheses is recommended for hemiarthroplasty to manage displaced femoral neck fracture with reduce of post-operative pain and better mobility.

Conclusion:

Functional outcome of hemiarthroplasty in elderly patients with fracture neck femur in terms of pain relief, mobility and restoration of function as near as possible to the pre-injury level was satisfactory in majority of patients with minimal morbidity.

Key words: Fracture, neck femur, elderly, hemiarthroplasty, functional result

Functional results of some studies with partial hemiarthroplasty in fracture neck femur in elderly patients versus internal fixation. Table (1-1)

Study name	Pain and Early mobility	Harris hip score	revision	mortality
Bhandari et al	Good results	HIGH score 80	Decreased rate	Low rate
Keating et al	Good results	Higher score 80	Decreased to 5%	Low rate
Abboud et al	Good results	Higher score 70	Decreased 8%	Low rate
Tidermark et al	Good results	Higher score 80	Decreased to 4%	Low rate

Raia et al	Good results	Higher score 80	Decreased to 5%	Low rate
Calder et al	Good results	Higher score	Decreased to 5%	Low rate

List of contents

Section No	Subject	Page. No
	Dedication	III
	Acknowledgment	IV
	Summary	V
1	Introduction	VII
2	Aim	1
3	Literatures' review	2-18
4	Conclusions	19-20
5	References	21-30

Chapter One

Introduction

Introduction

Fracture of femoral neck is common in clinical practice when elderly people are taken into consideration and the poor outcome in the form of morbidity and mortality in association with these fractures is also frequent ⁽¹⁾. Because of improvement in life expectancy, the incidence rate of these fractures has increased substantially and it is estimated that the incidence rate will triple by 2050 ⁽¹⁾. It has been reported that the mortality rate within one year of onset of these fractures is between 20 and 30% ⁽²⁾. These fractures by themselves are challenging to orthopedic surgeon and become more challenging because of the high rate of medical comorbidities in such elderly patients. It is well known that surgical intervention is the best approach in such fractures; however, the most fruitful surgical option is still controversial ⁽³⁾. Because of favorable advantages, such as early mobility, weight bearing that happens as soon as possible and better quality of life, hip replacement, whether total or partial, has gained nice reputation however 100 % consensus has not emerged in recent articles ⁽⁴⁾. Several studies have shown that the outcome of hemiarthroplasty is much better than internal fixation because of avoidance of avascular necrosis and non-union outcomes ⁽⁵⁻⁷⁾. Indeed unipolar hemiarthroplasty using Austin Moore prosthesis is still commonly practiced in developing countries for very limited or non-ambulatory elderly with relatively shorter life expectancy ⁽³⁾. Complications rates following unipolar and bipolar endoprosthesis have been cooperatively studies and the results have denied any significant difference in outcome between these two methods ⁽⁴⁾. The rarity of Iraqi literatures dealing with functional outcome of hemiarthroplasty among elderly patients in our community, this study was planned and carried out to make a clear insight on these outcomes.

Aim of this study: To evaluate the functional results after partial arthroplasty in compares to others modalities of management of fracture neck femur in elderly patients.

Chapter Two
Literatures Review
And discussion

2.1 Femoral Neck Fracture

Femoral neck fractures are intracapsular and typically occur in a bimodal age distribution, with most occurring in the elderly population. The incidence of femoral neck fractures increases with age. The patient's medical history and pre-injury status (i.e., prior hip pain, ambulatory status, functional and mental capacity) provide valuable information that may influence the treatment course (8,9).

2.2 Non-displaced Femoral Neck Fracture

Whether to manage nondisplaced femoral neck fractures nonsurgical or surgical is a topic of debate. Elderly patients with medical conditions that place them at high risk for anesthesia- and surgery-related complications can be treated nonsurgical. Nonambulatory patients and patients suffering from severe dementia who have minimal discomfort may also be treated nonsurgical (10).

Surgical fixation for nondisplaced fractures allows early patient mobilization and ensures that a nondisplaced fracture does not subsequently displace. Currently, there are no level I or II studies comparing nonsurgical with surgical management of nondisplaced femoral neck fractures (11). We evaluated two level III studies and one level IV study of nondisplaced femoral neck fracture. Hansen (12) performed a nonrandomized study involving 23 patients, 16 of whom were treated nonsurgically and 7 of whom were treated

surgically with sliding hip screws (SHSs). Nonunion occurred in 10 of 16 patients treated nonsurgical and in none of the surgically treated patients. Nine of 16 patients with a nonunion required revision surgery, whereas only 1 surgically treated patient required revision surgery. An 86% union rate was reported in one study of 170 consecutive patients with impacted femoral neck fractures who were treated with early mobilization and weight bearing (13). Patients older than age 70 years and in poor general health had the highest rate of secondary displacement. In a series of 1,400 patients, Parker et al (14) performed a cost-benefit analysis of various methods of treatment of hip fractures. The authors estimated a 30% 1-year mortality rate for patients whose nondisplaced subcapital fractures were treated nonsurgical and who had an uneventful union. For those patients with displaced subcapital fractures, the authors predicted a 90% 1-year mortality rate secondary to pneumonia, bedsores, and pulmonary emboli. Conn and Parker⁸ examined 375 patients with nondisplaced intracapsular fractures treated with internal fixation. The authors noted a nonunion rate of 6.4% and an osteonecrosis rate of 4.0%. Age, walking ability, degree of impaction evident on the anteroposterior radiograph, and angulations on the lateral radiograph were determined to be predictive of healing complications. In this study, the conversion rate to arthroplasty was 7.7%. Based on the available evidence, a recommendation cannot be made regarding the treatment of nondisplaced femoral neck fractures. The patient who is treated nonsurgical is not at risk of surgery related complications, including wound infections or complications associated with anesthesia. However, the nonunion rate is increased, as are complications associated with prolonged recumbence. Although further randomized trials would provide more

data, they may be difficult to conduct based on the modern standard of care and the limited indications for nonsurgical treatment.

2.3 Displaced Femoral Neck Fracture

The patient with a displaced femoral neck fracture is at significant risk for osteonecrosis and nonunion. Treatment options include closed reduction and internal fixation or ORIF (open reduction-internal fixation) with different constructs, hemiarthroplasty (unipolar and bipolar), and THA (total hip arthroplasty).

2.4 Internal Fixation

Many constructs have been used for internal fixation, including multiple screw fixations in a variety of configurations and SHSs (surgical helmet systems). In our attempt to determine whether a particular implant provides superior fixation, we assessed outcomes such as rates of nonunion and osteonecrosis, need for hardware removal, periprosthetic fracture, and implant failure. The following data are all from level I studies. A review of the Cochrane database revealed 28 randomized or quasirandomized trials of 5,547 patients with femoral neck fractures treated with 19 different pin and/or crew constructs in a variety of configurations (15). None of the implants had significantly superior results for outcomes related to fracture healing, osteonecrosis, wound infection, pain scores, reoperation rate, and use of walking aids, periprosthetic fracture, or mortality. Seven studies compared outcomes between SHSs and various cancellous screws. Four studies noted shorter surgical times with cancellous screws (average, 11 minutes) (16-18). One study reported surgical times to be equivalent between the fixation methods (19). In the SHS group, there was a tendency toward increased blood loss

(average, 84 mL), and deep wound infection was more common. Although the overall reoperation rate was equivalent between the groups, failure of fixation was lower in the SHS group. There was not a significant difference in mortality between the groups. Parker and Blundell¹⁴ conducted a meta-analysis of 25 randomized controlled clinical trials (RCTs) involving 4,925 patients with intracapsular fractures who were treated with a variety of implants. The review and analysis were focused on complications associated with fracture healing. No investigated device proved to be superior to any other in terms of nonunion or fracture displacement. There was limited evidence supporting screw fixation over smooth pins; however, this advantage was negated with the use of a hook at the end of the pin. No advantage was seen in using a side plate for fixation, and no significant evidence was found concerning the number of screws necessary for fixation. Based on the available evidence, there appear to be minimal differences between implants used for internal fixation of displaced femoral neck fractures (16, 18). These studies did not break down the data strictly based on age. Thus, it is not possible to recommend a particular implant for age-specific populations. The choice of implant should be based on surgeon familiarity and comfort level.

2.5 Internal Fixation versus Hemiarthroplasty

Multiple studies have been done on the outcomes of internal fixation of femoral neck fractures versus arthroplasty (eg, hemiarthroplasty, THA). The risk of osteonecrosis, nonunion, and revision following internal fixation of displaced intracapsular fractures must be

balanced against the potential complications following arthroplasty. A review of the Cochrane database produced 13 randomized or quasirandomized controlled trials with a total of 2,091 patients treated with either internal fixation or hemiarthroplasty. One clear limitation of this review is that different methods of internal fixation and different implants for arthroplasty were used. Outcomes assessed included surgical time, blood loss, wound infections, postoperative complications, and mortality (20, 21). Eight studies assessed the length of surgery, and all reported decreased surgical time for the patients treated with internal fixation (average, 22 minutes) (22). Additionally, the internal fixation group had a more favorable outcome in terms of blood loss, need for postoperative blood transfusions, and infection rates. No differences between the groups were found regarding mortality rates, pain, or mobility; however, there was a higher reoperation rate with internal fixation than with hemiarthroplasty (31% vs 8%; relative risk, 3.66). In an RCT of 100 patients conducted by Rödén et al (23) displaced femoral neck fractures were treated with either two von Bahr screws or a bipolar prosthesis. Inclusion criteria included age >70 years, no prior hip disease, ability to ambulate before injury, and no signs of senility. The duration of surgery was shorter and the blood loss less in the internal fixation group. This cohort did have a significant revision rate (34 of 53 patients). The prosthesis group was notable for a high dislocation rate (7 of 47 patients). No differences in patient mortality were noted at either 2- or 5-year follow-up. Parker et al (24) reported on 455 patients randomized to either internal fixation or hemiarthroplasty and found no differences in outcomes for pain, mobility, or mortality at 3-year follow-up. However, the authors did note a lower rate of revision in the hemiarthroplasty group (5%) than

in the group treated with internal fixation (40%). These results corroborated those of an earlier study by Parker and Pryor (25) of 208 patients treated with either internal fixation or hemiarthroplasty. An RCT of internal fixation versus hemiarthroplasty in the year 2001 was terminated early because of a 44% revision rate at 1 year in the internal fixation group (26). No revisions were needed in the hemiarthroplasty group. Rogmark et al (27) performed an RCT comparing internal fixation with hemiarthroplasty. At 2-year follow-up, patients who had undergone hemiarthroplasty had improved walking and stair climbing ability, and decreased pain levels. The available level II evidence seems to be consistent with the findings of level I studies, namely, that there is a higher revision rate for femoral neck fractures treated with internal fixation versus those managed with hemiarthroplasty. A meta-analysis by Lu-Yao et al (28) showed no difference in mortality in patients treated with internal fixation versus hemiarthroplasty, except for a non-statistically significant increase in the arthroplasty group in the first month (relative risk, 1.4). Nonunion developed in 33% of patients, and osteonecrosis in 16%, with reoperation rates for these complications ranging from

20% to 36%. In the arthroplasty group, the rates of reoperation for any cause ranged from 6% to 18%. The ideal treatment of displaced intracapsular fractures is not straightforward. The current data indicate that internal fixation of femoral neck fractures is associated with a greater number of significant problems (e.g., osteonecrosis, nonunion, revision) than is hemiarthroplasty. These risks outweigh the benefits of slightly shorter surgical times and marginally decreased blood loss. With similar mortality and pain scores,

hemiarthroplasty appears to be the better option for displaced femoral neck fractures. However, other factors critical in the decision-making process, such as age, were not considered in most of these studies.

Thus, we cannot make a definitive evidence-based judgment across all age groups and all circumstances as to the best treatment of displaced intracapsular hip fractures.

2.6 Cemented Versus Cementless Hemiarthroplasty

The first hip fracture endoprostheses were designed for cementless use, but cemented fixation has become the preferred technique with current femoral components. Numerous reports have documented improved outcomes with cemented implants. Emery et al (29) performed an RCT of 53 hemiarthroplasties. Twentyseven patients underwent cemented hemiarthroplasty, and 26 underwent cementless insertion. At a mean follow-up of 17 months, no statistically significant differences were noted between the groups with regard to postoperative complications, surgical time, estimated blood loss, or mortality. However, patients with cementless stems experienced a markedly higher level of hip pain and dependency on walking aids. In a Cochrane database review, Parker and Gurusamy (30) evaluated five trials with a total of 482 patients. Although there was no difference in complication or mortality rates, there was a higher rate of failure to regain preoperative mobility in the cementless prosthesis group. Additionally, patients with cementless prostheses noted higher pain scores at follow-up. The conclusion of this review was that cementing the prosthesis led to reduced pain postoperatively and

better mobility. However, in the trials, there was underreporting of outcomes in a small cohort of patients, which was felt to limit any definitive conclusions. In a systematic review of the literature (level II), Khan et al (31) found lower revision rates, less thigh pain, and better mobility in patients treated with cemented prostheses. There was no difference in general complications or mortality rates between the cemented and cementless groups. Lo et al (32) reviewed 451 displaced fractures of the femoral neck with at least 2-year follow-up and confirmed higher Harris hip scores and less thigh pain in the cemented group. The study reported longer average surgical times (20 minutes) and greater blood loss (160 mL) in the cemented group, but there was no significant difference in complication or mortality rates. A review of the SIGN database concerning cemented versus cementless stems corroborated the findings of other studies in that cemented prostheses resulted in improved mobility postoperatively and decreased pain. In addition, the use of cemented implants did not cause a higher rate of postoperative complications, increased surgical time, greater blood loss, or mortality. The SIGN database cited a study in which the authors concluded that cementless stems are associated with higher levels of thigh pain and lower overall hip scores (26). The recommendations by Dorr et al (33) suggested using a cemented prosthesis unless the patient exhibits cardiorespiratory compromise. In a level III study, Lennox and McLauchlan (34) treated 207 patients with either cemented or cementless hemiarthroplasty for displaced subcapital fractures of the femoral neck. Follow-up was done at an average of 19 months postoperatively. Mortality was higher in the cemented group than in the cementless group at 48 hours and at 3 months (4% versus 0%). Excluding the perioperative period, the number of postoperative days

until death was the same, suggesting that the use of cement may lead to higher mortality in the first 48 hours postoperatively. There were no differences in the overall complication rate, but patient satisfaction was higher in the cemented hemiarthroplasty group. The findings of equivalent complication rates and higher patient satisfaction with cemented hemiarthroplasty were recently confirmed by Singh and Deshmukh (28). However, in their review of 244 patients undergoing hemiarthroplasty, Foster et al (29) reported a higher rate of periprosthetic fracture in patients undergoing cementless versus cemented hemiarthroplasty (7% versus 0%). The authors noted that the patients in the cementless subgroup were significantly older; however, the ASA (American Society of Anesthesiologists) scores between the groups were comparable. Based on a review of the current evidence, we recommend using cemented prostheses when performing hemiarthroplasty to manage displaced femoral neck fractures. There is reasonable support for cemented fixation, with the decreased incidence of postoperative pain and better mobility.

2.7 Unipolar Versus Bipolar Hemiarthroplasty

Numerous studies have attempted to document possibly superior outcomes with the insertion of bipolar prostheses in patients with femoral neck fracture. Theoretically, because there is no prosthesis-prosthesis interface in a unipolar implant, the rate of acetabular wear should be reduced over time, and there should be a decreased incidence of pain and need for revision. A review of the Cochrane database included seven randomized or quasirandomized

trials involving 857 patients undergoing unipolar or bipolar hemiarthroplasty for femoral neck fracture (23). The results indicated no significant difference in acetabular wear, functional outcomes, length of surgery, blood loss, wound infections, or mortality. Raia et al (37) performed a prospective randomized trial comparing the efficacy of unipolar versus bipolar hemiarthroplasty in 115 patients older than age 65 years with displaced femoral neck fracture. Both types of prosthesis were cemented and implanted through a posterolateral approach. Patients were evaluated for quality of life and functional outcomes at 1 year postoperatively. The authors concluded that there was no difference in estimated blood loss, length of hospital stay, mortality rate, and number of dislocations, postoperative complications, or ambulatory status. Thus, the bipolar endoprosthesis provided no advantage in the treatment of displaced femoral neck fractures. Calder et al (38) performed a randomized prospective study in 250 patients aged 80 years and older to determine the rate of complications, patient satisfaction, Harris hip scores, degree of return to pre-injury state, and mortality. The only significant difference between the two groups involved return to pre-injury status, which was significantly greater ($P = 0.04$) following insertion of the unipolar prosthesis. The SIGN database states that a level II evidence comparing unipolar with bipolar prostheses. One study included a fluoroscopic evaluation of the bipolar prostheses (39). The radiologic data suggested that the majority of motion occurred at the outer articulation (acetabulum-prosthesis interface). There was little, if any, motion at the bipolar interface, which essentially served to convert the bipolar prosthesis to a unipolar device. The bipolar design was created to reduce acetabular wear as well as to minimize

pain and maximize mobility. The review concluded that these outcomes are related to patient activity level and duration of followup. The SIGN recommendation stated that bipolar hemiarthroplasty should not be performed in preference to unipolar hemiarthroplasty because there is limited evidence of clinical benefit with a bipolar prosthesis. Eiskjaer and Ostgård (40) reported on a total of 679 cases, which included 202 unipolar Austin Moore prostheses, 209 trunion-bearing Christiansen devices, and 268 Hastings bipolar hemiarthroplasty devices. The cumulative prosthesis survival was 90% at 5-year follow-up and 85% at 10-year follow-up. In contrast to other studies, significantly fewer failures were noted in the group undergoing cemented bipolar hemiarthroplasty. Yamagata et al (41) reported on 1,001 hemiarthroplasties (682 unipolar, 319 bipolar). The survivorship results of this level III study demonstrated a 13.7% revision rate at 8-year follow-up for bipolar components, compared with 22.9% for unipolar prostheses. In addition, patients undergoing bipolar hemiarthroplasty exhibited higher average Harris hip scores and lower acetabular erosion rates compared with those managed with the unipolar devices. In a level IV study, Haidukewych et al (42) reviewed the results and survivorship of 212 cemented bipolar hemiarthroplasties in 205 patients (average age, 79 years). The authors reported 10-year survivorship free of revision for any reason of 93.6%. This rate increased to 95.9% when factors other than mechanical failure (i.e., aseptic loosening, acetabular wear) were excluded. The authors concluded that cemented bipolar hemiarthroplasty was associated with excellent component survivorship in elderly patients. They documented a low complication rate (11%) and satisfactory pain relief (96%). Given

the variable conclusions from numerous levels I through IV studies concerning hemiarthroplasty, absolute recommendations cannot be made concerning the type of implant to be used. There is limited evidence to support the use of a bipolar prosthesis over unipolar designs. Although there is a theoretical design advantage with bipolar implants, these advantages have yet to be confirmed in clinical studies. Additional large-scale RCTs are needed to definitively answer this question.

2.8 Surgical Approach

Surgical approach reportedly affects the incidence of dislocations and infections, duration of surgery, and blood loss. Most frequently, an anterolateral or a posterior approach is performed. The question is whether there is an ideal surgical approach that minimizes complications and causes the least possible morbidity. No level I study has specifically evaluated the surgical approach. A review of the Cochrane database identified only one RCT of 114 patients comparing surgical approaches for hemiarthroplasty (level II). Sikorski and Barrington (43) followed for 2 years 57 patients who had undergone either an anterolateral or a posterior approach for displaced subcapital femoral fracture. A cemented Thompson prosthesis was used for all patients. The rates of dislocation, prosthesis loosening, acetabular protrusion, wound infection, and revision were similar between the groups. Of note, medical complications, including pneumonia, congestive heart failure, and urinary tract infections, were higher in the posterior approach group, as was postoperative mortality (25% vs. 42% at 2-year follow-up).

The postoperative protocol followed at that time (i.e., prolonged bed rest to prevent posterior dislocation) may have contributed to these findings. Additionally, selection bias and poor, somewhat inconsistent reporting of results may have compromised the integrity of the study. Keene and Parker (44) conducted a prospective study of 531 patients who underwent hemiarthroplasty with either an anterior or a posterior approach. The anterolateral approach was associated with increased surgical time (8 minutes longer), blood loss (54 mL), and superficial infection (6% versus 2.6%). However, the report also indicated that the posterior approach was associated with a higher dislocation rate (4.3% versus 1.7%) and more thromboembolic complications (9.2% versus 1.3%). There was no difference in hospital stay or mortality, and the authors suggested that surgeon comfort with the approach should dictate the exposure used. In their level IV case series of 1,812 bipolar hemiarthroplasties, Sierra et al (45) found no significant differences in their comparison of dislocation rates between anterolateral, posterolateral, and transtrochanteric approaches. The authors noted a total of only 32 dislocations, half of which occurred during the first 6 months postoperatively. In 2004, Varley and Parker (46) performed a systematic literature review of dislocations and surgical approach during a 40-year period. They found that the rate of dislocation with a posterior approach was 5.1%, compared with 2.4% for an anterior approach. There is a lack of strong evidence to advocate one particular surgical approach for hemiarthroplasty. In the absence of data from well-designed RCTs, the choice of surgical approach will be based primarily on the surgeon's clinical assessment of each patient's needs and the surgeon's surgical experience.

2.9 Internal Fixation versus Total Hip Arthroplasty

Indications for THA following a displaced femoral neck fracture have included the presence of preexisting osteoarthritis, rheumatoid arthritis, and degenerative joint disease secondary to Paget's disease. However, more recently there has been increased enthusiasm for primary THA for managing displaced femoral neck fractures. Tidermark et al (40) conducted a prospective CT of 102 patients (mean age, 80 years) with displaced femoral neck fractures treated with either internal fixation or THA. Outcomes measures included hip function, quality of life, complications, and revision surgery. At 2-year follow-up, the complication rate (36% versus 4%, $P < 0.001$) and revision rate (42% versus 4%, $P < 0.001$) were significantly higher in the internal fixation group than in patients treated with THA. Hip function in terms of quality of life ($P < 0.05$), comfort ($P < 0.005$), motion ($P < 0.05$), and walking ability ($P < 0.05$) were all significantly better in this group of independent, cognitively intact patients treated with THA. At 4-year follow-up, the same investigators reported that the incidence of complications and revisions in the internal fixation group had increased but that no additional complications were reported, and no revisions were required in the arthroplasty group (41). Similar results were seen in a larger RCT conducted by Rogmark et al (49) who noted improved pain scores ($P < 0.05$) and walking ability ($P < 0.05$) in the THA group. The mortality rate at 2-year follow-up was 21% for both groups, with a higher mortality rate among men (33% versus 18%). Johansson et al (50) also found an increased rate of complications at 1-year follow-up in patients with intact cognition who underwent

internal fixation. Patients with compromised mental status had a higher rate of complications and a higher mortality rate at 2 years following ORIF (57.7% versus 12.7%). Ravikumar and Marsh (51) performed an RCT of 290 patients older than age 65 years, comparing internal fixation, hemiarthroplasty, and THA. At 13-year follow-up, revision rates were the lowest (6.75%) and Harris hip scores were the highest (80) in the patients who had undergone THA. The internal fixation and hemiarthroplasty groups had revision rates of 33% and 24%, respectively. Skinner et al (52) also randomized 278 patients to ORIF, hemiarthroplasty, or THA for displaced femoral neck fractures. They showed equivalent mortality at 1 year postoperatively (25%). The internal fixation group exhibited the highest revision rate (25%). Pain relief and mobility were best in the THA group. Bhandari et al (53) conducted a metaanalysis of all RCTs reported over a 33-year period, comparing internal fixation and arthroplasty (ie, hemiarthroplasty, bipolar arthroplasty, THA). Cumulative data showed a decreased rate of revision surgery in the arthroplasty group and an increased risk of infection. The relative risk of mortality in the arthroplasty group was higher during the first 4 months postoperatively but was no longer evident at 1-year follow-up. The most recent multicenter RCT comparing internal fixation, hemiarthroplasty, and THA in cognitively intact patients was performed by Keating et al⁴⁷ (level II). At 2-year follow-up, revision surgery was required in 39% of the internal fixation group, 5% of the hemiarthroplasty group, and 9% of the THA group. Hip scores and quality of life measurements were significantly greater in the THA group than in the other two groups. Economic analysis revealed that internal fixation was least costly to perform acutely but was most costly after including the cost of

additional treatment of complications. As the popularity of treating femoral neck fractures with THA has increased, outcomes have been compared with those of patients undergoing THA for degenerative conditions. A retrospective study of 60 patients by Abboud et al (55) showed no difference in outcomes for patients undergoing THA for femoral neck fractures versus those undergoing THA for osteoarthritis. Harris hip scores, perioperative morbidity, and mortality were equivalent for both groups. This is in contrast to earlier studies that showed increased rates of dislocation in patients undergoing primary THA for femoral neck fractures. Current level I and II evidence indicates that as the index procedure for a femoral neck fracture, THA leads to better outcomes than internal fixation, as measured by hip function scores and a decreased rate of revision surgery. This option should be strongly considered for the healthy, cognitively intact patient. Ultimately, more RCTs are needed to further clarify the risks and benefits of each procedure for clearly defined patient groups.

Chapter Three

Conclusions

Conclusion

Functional outcome of hemiarthroplasty in elderly patients with fracture neck femur in terms of pain relief, mobility and restoration of function as near as possible to the pre-injury level was satisfactory in majority of cases with minimal morbidity

Careful patient's selection for hemiarthroplasty is vital and may decrease the incidence of complications and ameliorate the outcomes in the treatment of displaced intracapsular femoral neck fractures.

Primary cemented hemiarthroplasty rather than uncemented endoprosthesis may be favored in the treatment of displaced intracapsular fracture neck femur in elderly patients above 60 years.

References

1. Schmidt AH, Swiontkowski MF. Femoral neck fractures. *Orthop Clin North Am.* 2002;3:97–111.
2. Daniel M, Mohammed S, Francis A, William Y, Joseph K, Cornilius E. Early result of hemiarthroplasty in elderly patients with fracture neck of femur. *Niger Med J.* 2015;56(1):64-8.
3. Nather A, Seow CS, Iau P, Chan A. Mortality and morbidity for elderly patients with fractured neck of femur treated by hemiarthroplasty. *Injury.* 1995;26:187–90.
4. Calder SJ, Anderson GH, Jagger C, Harper WM, Gregg PJ. Unipolar or bipolar prosthesis in displaced intracapsular hip fractures in octogenarians: A randomised prospective study. *J Bone Joint Surg Br.* 1996;78:391–4.
5. Rogmark C, Carlsson A, Johnell O, Sernbo I. Primary hemiarthroplasty in old patients with displaced femoral neck fracture: A 1-year follow-up of 103 patients aged 80 years or more. *Acta Orthop Scand.* 2002;73:605–10.
6. Roden M, Schon M, Freden H. Treatment of displaced femoral neck fractures: A randomized minimum 5-year follow-up study of screws and bipolar hemiprosthesis in 100 patients. *Acta Orthop Scand.* 2003;74:42–4.
7. Parker MJ, Khan RJ, Crawford J, Pryor GA. Hemiarthroplasty versus internal fixation for displaced

intracapsular hip fractures in the elderly. A randomized trial of 455 patients. *J Bone Joint Surg Br*. 2002;84:1150–5.

8. Stevens JA, Olson S: Reducing falls and resulting hip fractures among older women. *MMWR Recomm Rep* 2000;49(RR-2):3-12.

9. USCensus Bureau:US Interim Projections by Age, Sex, Race, and Hispanic Origin. Washington, DC: US Census Bureau, March 18, 2004. Available at: <http://www.census.gov/ipc/www/usinterimproj/natprojtab02a.pdf>.

10. Kannus P, Parkkari J, Sievänen H, Heinonen A, Vuori I, Järvinen M: Epidemiology of hip fractures. *Bone* 1996;18(1 suppl):57S-63S.

11. Zuckerman JD: Hip fracture. *N Engl J Med* 1996;334:1519-1525.

12. Hansen F: Conservative vs surgical treatment of impacted, subcapital fractures of the femoral neck. *Acta Orthop Scand* 1994;256:9.

13. Raaymakers EL, Marti RK: Nonoperative treatment of impacted femoral neck fractures: A prospective study of 170 cases. *J Bone Joint Surg Br* 1991;73:950-954.

14. Parker MJ, Myles JW, Anand JK, Drewett R: Cost-benefit analysis of hip fracture treatment. *J Bone Joint Surg Br* 1992;74:261-264.

15. Conn KS, Parker MJ: Undisplaced intracapsular hip fractures: Results of internal fixation in 375 patients. *Clin Orthop Relat Res* 2004;421:249-254.
16. Parker MJ, Stockton G: Internal fixation implants for intracapsular proximal femoral fractures in adults. *Cochrane Database Syst Rev* 2001;4: CD001467.
17. Benterud JG, Husby T, Nordsletten L, Alho A: Fixation of displaced femoral neck fractures with a sliding screw plate and a cancellous screw or two Olmed screws: A prospective, randomized study of 225 elderly patients with a 3-year follow-up. *Ann Chir Gynaecol* 1997;86:338-342.
18. Madsen F, Linde F, Andersen E, Birke H, Hvass I, Poulsen TD: Fixation of displaced femoral neck fractures: A comparison between sliding screw plate and four cancellous bone screws. *Acta Orthop Scand* 1987;58:212-216.
19. Paus A, Gjengedal E, Hareide A, Jorgensen JJ: Dislocated fractures of the femoral neck treated with von Bahr screws or hip compression screw: Results of a prospective, randomized study. *J Oslo City Hosp* 1986;36:55- 61.
20. Kuokkanen H, Korkala O, Antti-Poika I, Tolonen J, Lehtimäki MY, Silvennoinen T: Three cancellous bone screws versus a screw-angle plate in the treatment of Garden I and II fractures of the femoral neck. *Acta Orthop Belg* 1991;57:53-57.
21. Parker MJ, Blundell C: Choice of implant for internal fixation of femoral neck fractures: Meta-analysis of 25

randomised trials including 4,925 patients. Acta Orthop Scand 1998;69: 138-143.

22. Masson M, Parker MJ, Fleischer S: Internal fixation versus arthroplasty for intracapsular proximal femoral fractures in adults. Cochrane Database Syst Rev 2003;2:CD001708.

23. Rödén M, Schön M, Fredin H: Treatment of displaced femoral neck fractures: A randomized minimum 5-year follow-up study of screws and bipolar hemiprotheses in 100 patients. Acta Orthop Scand 2003;74:42-44.

24. Parker MJ, Khan RJ, Crawford J, Pryor GA: Hemiarthroplasty versus internal fixation for displaced intracapsular hip fractures in the elderly: A randomized trial of 455 patients. J Bone Joint Surg Br 2002;84:1150-1155.

25. Parker MJ, Pryor GA: Internal fixation or arthroplasty for displaced cervical hip fractures in the elderly: A randomized controlled trial of 208 patients. Acta Orthop Scand 2000;71: 440-446.

26. Puolakka TJ, Laine HJ, Tarvainen T, Aho H: Thompson hemiarthroplasty is superior to Ullevaal screws in treating displaced femoral neck fractures in patients over 75 years: A prospective randomized study with two-year follow-up. Ann Chir Gynaecol 2001; 90:225-228.

27. Rogmark C, Carlsson A, Johnell O, Sembol: Costs of internal fixation and arthroplasty for displaced femoral neck fractures: A randomized study of 68 patients. Acta Orthop Scand 2003;74:293-298.

28. Lu-Yao GL, Keller RB, Littenberg B, Wennberg JE: Outcomes after displaced fractures of the femoral neck: A meta-analysis of one hundred and six published reports. *J Bone Joint Surg Am* 1994;76:15-25.
29. Emery RJ, Broughton NS, Desai K, Bulstrode CJ, Thomas TL: Bipolar hemiarthroplasty for subcapital fracture of the femoral neck: A prospective randomised trial of cemented Thompson and uncemented Moore stems. *J Bone Joint Surg Br* 1991;73: 322-324.
30. Parker MJ, Gurusamy K: Arthroplasties (with and without bone cement) for proximal femoral fractures in adults. *Cochrane Database Syst Rev* 2004;2:CD001706.
31. Khan RJ, MacDowell A, Crossman P, Keene GS: Cemented or uncemented hemiarthroplasty for displaced intracapsular fractures of the hip: A systematic review. *Injury* 2002;33:13-17.
32. Lo WH, Chen WM, Huang CK, Chen TH, Chiu FY, Chen CM: Bateman bipolar hemiarthroplasty for displaced intracapsular femoral neck fractures: Uncemented versus cemented. *Clin Orthop Relat Res* 1994;302:75-82.
33. Dorr LD, Glousman R, Hoy AL, Vanis R, Chandler R: Treatment of femoral neck fractures with total hip replacement versus cemented and noncemented hemiarthroplasty. *J Arthroplasty* 1986;1:21-28.

34. Lennox IA,McLauchlan J: Comparing the mortality and morbidity of cemented and uncemented hemiarthroplasties. *Injury* 1993;24:185-186.
35. Singh GK, Deshmukh RG: Uncemented Austin-Moore and cemented Thompson unipolar hemiarthroplasty for displaced fracture neck of femur: Comparison of complications and patient satisfaction. *Injury* 2006;37: 169-174.
36. Foster AP, Thompson NW, Wong J, Charlwood AP: Periprosthetic femoral fractures: A comparison between cemented and uncemented hemiarthroplasties. *Injury* 2005;36:424-429.
37. Raia FJ, Chapman CB, Herrera MF, Schweppe MW, Michelsen CB, Rosenwasser MP: Unipolar or bipolar hemiarthroplasty for femoral neck fractures in the elderly? *Clin Orthop Relat Res* 2003;414:259-265.
38. Calder SJ, Anderson GH, Jagger C, Harper WM, Gregg PJ: Unipolar or bipolar prosthesis for displaced intracapsular hip fracture in octogenarians: A randomised prospective study. *J Bone Joint Surg Br* 1996;78:391-394.
39. Eiskjaer S, Gelineck J, Søballe K: Fractures of the femoral neck treated with cemented bipolar hemiarthroplasty. *Orthopedics* 1989;12:1545-1550.
40. Eiskjaer S, Ostgård SE: Survivorship analysis of hemiarthroplasties. *Clin Orthop Relat Res* 1993;286:206-211.

41. Yamagata M, Chao EY, Ilstrup DM, Melton LJ III, Coventry MB, Stauffer RN: Fixed-head and bipolar hip endoprostheses: A retrospective clinical and roentgenographic study. *J Arthroplasty* 1987;2:327-341.
42. Haidukewych GJ, Israel TA, Berry DJ: Long-term survivorship of cemented bipolar hemiarthroplasty for fracture of the femoral neck. *Clin Orthop Relat Res* 2002;403:118-126.
43. Sikorski JM, Barrington R: Internal fixation versus hemiarthroplasty for the displaced subcapital fracture of the femur: A prospective randomized study. *J Bone Joint Surg Br* 1981;63: 357-361.
44. Keene GS, Parker MJ: Hemiarthroplasty of the hip: The anterior or posterior approach? A comparison of surgical approaches. *Injury* 1993;24: 611-613.
45. Sierra RJ, Schleck CD, Cabanela ME: Dislocation of bipolar hemiarthroplasty: Rate, contributing factors, and outcome. *Clin Orthop Relat Res* 2006;442:230-238.
46. Varley J, Parker MJ: Stability of hip hemiarthroplasties. *Int Orthop* 2004; 28:274-277.
47. Tidermark J, Ponzer S, Svensson O, Söderqvist A, Törnkvist H: Internal fixation compared with total hip replacement for displaced femoral neck fractures in the elderly: A randomised, controlled trial. *J Bone Joint Surg Br* 2003;85:380-388.

48. Blomfeldt R, Törnkvist H, Ponzer S, Söderqvist A, Tidermark J: Comparison of internal fixation with total hip replacement for displaced femoral neck fractures: Randomized, controlled trial performed at four years. *J Bone Joint Surg Am* 2005;87:1680-1688.
49. Rogmark C, Carlsson A, Johnell O, Sernbo I: A prospective randomized trial of internal fixation versus arthroplasty for displaced fractures of the neck of the femur: Functional outcome for 450 patients at two years. *J Bone Joint Surg Br* 2002;84:183-188.
50. Johansson T, Jacobsson SA, Ivarsson I, Knutsson A, Wahlström O: Internal fixation versus total hip arthroplasty in the treatment of displaced femoral neck fractures: A prospective randomized study of 100 hips. *Acta Orthop Scand* 2000;71:597-602.
51. Ravikumar KJ, Marsh G: Internal fixation versus hemiarthroplasty versus total hip arthroplasty for displaced subcapital fractures of femur: 13 year results of a prospective randomized study. *Injury* 2000;31:793-797.
52. Skinner P, Riley D, Ellery J, Beaumont A, Coumine R, Shafiqhian B: Displaced subcapital fractures of the femur: A prospective randomized comparison of internal fixation, hemiarthroplasty and total hip replacement. *Injury* 1989;20:291-293.
53. Bhandari M, Devereaux PJ, Swiontkowski MF, et al: Internal fixation compared with arthroplasty for displaced

fractures of the femoral neck: A meta-analysis. *J Bone Joint Surg Am* 2003;85:1673-1681.

54. Keating JF, Grant A, Masson M, Scott NW, Forbes JF: Randomized comparison of reduction and fixation, bipolar hemiarthroplasty, and total hip arthroplasty: Treatment of displaced intracapsular hip fractures in healthy older patients. *J Bone Joint Surg Am* 2006;88:249-260.

55. Abboud JA, Patel RV, Booth RE Jr, Azarian DG: Outcomes of total hip arthroplasty are similar for patients with displaced femoral neck fractures and osteoarthritis. *Clin Orthop Relat Res* 2004;421:151-154.

